

## Effect of zirconium chelate on cadmium stressed wheat (*Triticum aestivum* L. MV.20) seedlings

Marietta Fodor\*, Attila Hegedűs, Éva Stefanovits-Bányai

Department of Applied Chemistry, Corvinus University of Budapest, Budapest, Hungary

**ABSTRACT** Wheat seedlings were grown hydroponically, for the heavy metal treatment 200  $\mu\text{M}$  Cd acetate was applied. Nutrient solutions were also supplemented with a water-soluble zirconium ascorbate complex (Zr-ASC) at several concentrations. Our results indicated that wheat seedlings have accumulated great amounts of zirconium and cadmium in their roots and shoots. Zirconium uptake was increased under Cd stress; however, its translocation to the leaves was mitigated. Increased activities of guaiacol peroxidase (POD), ascorbic acid peroxidase (APX) and glutathione reductase (GR) reflected a Cd induced oxidative challenge in plant cells. Cadmium induced heavy metal stress symptoms could be alleviated by 10 and 33  $\mu\text{M}$  Zr-ASC, which was also confirmed by the reduced activities of all the tested antioxidant enzymes.

Acta Biol Szeged 49(1-2):93-94 (2005)

### KEY WORDS

Zirconium  
Cadmium  
wheat  
oxidative stress  
peroxidases  
glutathione reductase

Zirconium is the 20<sup>th</sup> most common element in the earth's crust, its total content in soils is deemed to be between 30 and 2000  $\text{mg kg}^{-1}$  dry soil. Zr is present in soil in amounts higher than nickel, zinc, copper or lead. Due to the changing environmental conditions or sources of contamination, water soluble Zr may occur in ecosystems. Water solubility means that Zr uptake by the roots becomes possible and in this way it may enter the food chain. This may have unknown influences on the domestic animals or even on the human body.

Our knowledge on the effects of Zr on living organisms is especially restricted. However, in a previous study we could detect favourable physiological effects on wheat seedlings growing in 10-55  $\mu\text{M}$  Zr-ASC (Fodor et al. 2005). We cannot rule out the possibility that this beneficial effect may even be more intensive in plants grown under severe abiotic stress conditions, as it was shown in case of titanium ascorbate, a very similar compound to Zr-ASC, in some crops (Alcaraz-Lopez et al. 2003).

Our experiments were carried out to survey that 10, 33 and 55  $\mu\text{M}$  Zr-ASC treatment could have beneficial effects on Cd-treated plants or not.

### Materials and Methods

For each treatment 3 g wheat caryopses (*Triticum aestivum* L. MV. 20) were imbibed for 24 hours and germinated in hydroponics of Knopp solution (Suba 1978), which was supplemented with 200  $\mu\text{M}$  Cd acetate. K-ASC and different concentrations of Zr-ASC (10, 33 55  $\mu\text{M}$  Zr) were used for Zr treatment. Zirconium was provided as Zr-ascorbate chelate. It was obtained from zirconyl chloride ( $\text{ZrOCl}_2 \cdot 8 \text{H}_2\text{O}$ ) and L-ascorbic acid (Fodor et al. 2003). Plants were kept in com-

plete darkness during the first 5 days of their development, and then grown further at room temperature (25°C) under natural irradiation. Nine-day-old seedlings were rinsed with distilled water, dried, and prepared for analysis.

Cd content was measured by ICP-OES. Plant material was dehydrated at 105°C for 24 h and 0.2 g of powdered dry material was dissolved in 2 ml of  $\text{HNO}_3 : \text{H}_2\text{O}_2$  mixture (1:1, v/v). Decomposition of plant tissues was completed by incubation at high pressure and temperature. Arsenazo III was used as a reagent to determine Zr content of the filtered solutions (Savvin 1961). The measurement was carried out by a PC-controlled GBC 916 UV/VIS spectrophotometer at  $\lambda = 665 \text{ nm}$  ( $\epsilon = 120 \text{ mmol}^{-1} \text{ cm}^{-1}$ ). Enzyme extraction and activity determination of guaiacol peroxidase (POD), ascorbate peroxidase (APX) and glutathione reductase (GR) was carried out as described by Fodor et al. (2005).

Three independent experiments were conducted and plants growing in 200  $\mu\text{M}$  cadmium acetate were taken as control. Statistical analyses were performed by Microsoft Excel software ( $P \leq 0.05$ ).


### Results

#### Zr content

Cadmium did not inhibit the Zr uptake of plants, in fact it was enhanced: 1200  $\mu\text{g/g}$  dry matter in case of 55  $\mu\text{M}$  Zr in contrast to the 430  $\mu\text{g/g}$  dry matter without Zr.

Zr translocation from roots to leaves was also considerably affected. Leaves of control plants contained 120  $\mu\text{g Zr/g}$  dry matter, while 55  $\mu\text{M}$  Zr treatment caused a marked decrease as leaves contained only 30  $\mu\text{g Zr/g}$  dry matter. Zirconium could not be detected in leaves of plants growing in Knopp solution.

\*Corresponding author. E-mail: marietta.fodor@uni-corvinus.hu



**Figure 1.** Zr-ASC induced changes in the POD activity in leaves and roots of Cd-stressed plants (\* =  $P < 0.05$ ).

### Determination of enzyme activities

Leaves and roots of control plants growing in Cd-containing nutrient solution performed increased POD activities compared to those growing in Cd-free solutions (Fig. 1). 10 and 33  $\mu\text{M}$  Zr-ASC treatment significantly reduced the POD activity in roots of plants. Their values have practically reached the level of POD activity measured in plants growing in Cd-free nutrient solutions. In leaves no alterations attributable to the effect of zirconium could be observed.

Cd treatment has considerably increased the APX activity in roots and leaves, while the additional Zr-ASC treatment in 10 and 33  $\mu\text{M}$  concentrations resulted in a significant reduction. 55  $\mu\text{M}$  Zr-ASC, however, markedly increased the APX activity in roots.

Cd treatment also induced elevated GR activities. In roots, a slight but not significant decrease was measured in plants treated by Zr, however, in leaves, 10 and 33  $\mu\text{M}$  Zr-ASC significantly decreased the GR activity to the level observed in plants grown in the K-ASC solution.

### Discussion

Cadmium does not generate active oxygen species directly, but by modifying some of the enzymes' activity, it induces oxidative stress (Hegedűs et al. 2001). Cd treatment induced considerable increases in all enzyme activities (POD, APX, GR) both in roots and leaves, which indicates that plants

suffered oxidative stress. In roots, 10 and 33  $\mu\text{M}$  Zr-ASC significantly lowered the POD and APX activities, indicating that small amounts of Zr may alleviate Cd induced oxidative stress effects. Similar results were presented for titanium ascorbate by Hegedűs et al. (2002) in case of barley seedlings, and by Koczka et al. (2001) in *Ginkgo biloba* L.

There was no remarkable increase in POD activity, when plants were grown in 200  $\mu\text{M}$  Cd-containing nutrient solution supplemented with 10–55  $\mu\text{M}$  Zr-ASC. 33 and 55  $\mu\text{M}$  Zr induced a slight but not significant elevation in POD activity, which can be attributed to a joint effect produced by the two different metals.

Activity increases of APX and GR were considerably restricted in leaves of plants treated by 10 and 33  $\mu\text{M}$  Zr-ASC compared to that of the Cd-treated control plants. This may reflect that Cd induced oxidative stress some way could be moderated by Zr-ASC supplementation in a concentration range of 10 and 33  $\mu\text{M}$ , and consequently, the activities of antioxidant enzymes were not as much elevated.

Enzyme activities in case of 55  $\mu\text{M}$  Zr treatment were similar to those of the Cd-stressed control plants, which indicates a concentration threshold, over which no protective effects can be demonstrated.

### Acknowledgements

This research was supported by the Hungarian National Scientific Research Fund (OTKA T043063).

### References

- Alcaraz-Lopez C, Botia M, Alcaraz CF, Riquelme F (2003) Effects of foliar sprays containing calcium, magnesium and titanium on plum (*Prunus domestica* L.) fruit quality. *J Plant Physiol* 160:1441-1446.
- Chance B, Maehly AC (1955) Assay of catalases and peroxidases. *Methods Enzymol* 2:764-817.
- Fodor M, Hegóczki J, Vereczkey G (2003) The effects of zirconium, a less known microelement, on basic fermentation characteristics and protein composition of *Saccharomyces cerevisiae*. *Acta Aliment* 32:353-362.
- Fodor M, Hegedűs A, Stefanovits-Bányai É (2005) Zirconium induced physiological alterations in wheat seedlings. *Biol Plantarum* (in press).
- Hegedűs A, Tóth E, Vajda K, Hauptmann G, Stefanovits-Bányai É, Erdei S (2002) Titán-askorbát kezelés hatása a növények antioxidáns védekező rendszerére. *Proc. The 9<sup>th</sup> Symposium on Analytical and Environmental Problems, Szeged*. 65-69.
- Koczka N, Orlóci L, Dimény J, Stefanovits-Bányai É (2001) Effect of cadmium stress on peroxidase activity of *Ginkgo biloba* L. *Proc. The 9<sup>th</sup> International Trace Element Symposium Budapest*, pp.159-165.
- Nakano Y, Asada K (1981) Hydrogen peroxide is scavenged by ascorbate specific peroxidase in spinach chloroplasts. *Plant Cell Physiol* 22:867-880.
- Savvin SB (1961) Analytical use of Arsenazo III. Determination of thorium, zirconium, uranium and rare earth elements. *Talanta* 8:673-685.
- Suba J (1978) *Növényélettani gyakorlatok*. Tankönyvkiadó, Budapest.